


**Amendments to the Claims**

This listing of the claims will replace all prior versions and listing of claims in this application.

**Listing of Claims**

1. (Previously Amended) A method for increasing the endoparasitocidal action of cyclic depsipeptides consisting of amino acids and hydroxycarboxylic acids as ring units and having 24 ring atoms, comprising contacting endoparasites with said cyclic depsipeptides in combination with piperazines.
2. (Previously Amended) An endoparasitocidal composition that contains piperazines together with cyclic depsipeptides consisting of amino acids and hydrocarboxylic acids as ring units and having 24 ring atoms.
- 3-6. (Withdrawn)
7. (Previously Amended) The method of claim 1 wherein said piperazines correspond to the formula (X),




(X)

in which

R<sup>13</sup> and R<sup>14</sup> independently of one another represent identical or different substituents of the group hydrogen, in each case optionally substituted alkyl, cycloalkyl, aryl, heteroaryl, and -CONR<sup>15</sup>R<sup>16</sup> or -CSNR<sup>15</sup>R<sup>16</sup>, in which

R<sup>15</sup> and R<sup>16</sup> independently of one another represent identical or different substituents of the group hydrogen, in each case optionally substituted alkyl or cycloalkyl.

8. (Currently Amended) The method of claim 1, wherein the piperazines correspond to the formula (X),




(X)

in which

R<sup>13</sup> and R<sup>14</sup> independently of one another represent identical or different substituents of the group hydrogen, in each case optionally substituted C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl, and -CONR<sup>15</sup>R<sup>16</sup> or -CSNR<sup>15</sup>R<sup>16</sup>, in which

R<sup>15</sup> and R<sup>16</sup> independently of one another represent identical or different substituents of the group hydrogen, in each case optionally substituted C<sub>1</sub>-C<sub>6</sub>-alkyl or C<sub>3</sub>-C<sub>8</sub>-cycloalkyl.

9. (Previously Amended) The method of claim 1, wherein the piperazines correspond to the formula (X)




(X)

in which

$R^{13}$  and  $R^{14}$  independently of one another represent identical or different substituents of the group hydrogen, in each case optionally substituted  $C_1$ - $C_4$ -alkyl,  $C_6$ -cycloalkyl, and  $-\text{CONR}^{15}\text{R}^{16}$  or  $-\text{CSNR}^{15}\text{R}^{16}$ , in which

$R^{15}$  and  $R^{16}$  independently of one another represent identical or different substituents of the group hydrogen, in each case optionally substituted  $C_1$ - $C_4$ -alkyl or  $C_6$ -cycloalkyl.

10. (Currently Amended) The composition as claimed in claim 2, wherein the cyclic depsipeptides correspond to the formula (I)



in which


$R^1$ ,  $R^2$ ,  $R^{11}$  and  $R^{12}$  independently of one another represent  $C_{1-8}$ -alkyl,  $C_{1-8}$ -halogenoalkyl,  $C_{3-6}$ -cycloalkyl, aralkyl, aryl,

$R^3$ ,  $R^5$ ,  $R^7$ ,  $R^9$  independently of one another represents hydrogen or straight chain or branched  $C_{1-8}$ -alkyl, which can optionally be substituted by hydroxyl,  $C_{1-4}$ -alkoxy, carboxyl ( $-\text{COOH}$ ), carboxamide ( $-\text{OCONH}_2$ ), imidazolyl, indolyl, guanidino,  $-\text{SH}$  or  $C_{1-4}$ -alkylthio and further represents aryl or aralkyl which can be substituted by halogen, hydroxyl,  $C_{1-4}$ -alkyl,  $C_{1-4}$ -alkoxy,

$R^4$ ,  $R^6$ ,  $R^8$ ,  $R^{10}$  independently of one another represent hydrogen, straight-chain  $C_{1-5}$ -alkyl,  $C_{2-6}$ -alkenyl,  $C_{3-7}$ -cycloalkyl, each of which can optionally be substituted by hydroxyl,  $C_{1-4}$ -alkoxy, carboxyl, carboxamide, imidazolyl, indolyl, guanidine, SH or  $C_{1-4}$ -alkylthio, and represent aryl or aralkyl which can be substituted by halogen, hydroxyl,  $C_{1-4}$ -alkyl,  $C_{1-4}$ -alkoxy,

and their optical isomers and racemates,

and the piperazines correspond to the formula (X),




(X)

in which


$R^{13}$  and  $R^{14}$  independently of one another represent identical or different substituents of the group hydrogen, in each case optionally substituted alkyl, cycloalkyl, aryl, **heterearyl-heteroaryl**, and  $-\text{CONR}^{15}\text{R}^{16}$  or  $-\text{CSNR}^{15}\text{R}^{16}$ , in which

$R^{15}$  and  $R^{16}$  independently of one another represent identical or different substituents of the group hydrogen, in each case optionally substituted alkyl or cycloalkyl.

11. (New) An endoparasitocidal composition that contains piperazines together with cyclic depsipeptides consisting of amino acids and hydrocarboxylic acids as ring units and having 24 ring atoms, wherein the cyclic depsipeptides correspond to the formula



and wherein the piperazines correspond to the formula (X),




(X)

in which


$R^{13}$  and  $R^{14}$  independently of one another represent identical or different substituents of the group hydrogen, in each case optionally substituted alkyl, cycloalkyl, aryl, heteroaryl, and  $-\text{CON} R^{15} R^{16}$  or  $-\text{CSN} R^{15} R^{16}$ , in which

$R^{15}$  and  $R^{16}$  independently of one another represent identical or different substituents of the group hydrogen, in each case optionally substituted alkyl or cycloalkyl.

12. (New) An endoparasitocidal composition that contains piperazines together with cyclic depsipeptides consisting of amino acids and hydrocarboxylic acids as ring units and having 24 ring atoms, wherein the cyclic depsipeptides correspond to the formula



and wherein the piperazines correspond to the formula (X),



(X)

in which

$R^{13}$  and  $R^{14}$  independently of one another represent identical or different substituents of the group hydrogen, in each case optionally substituted alkyl, cycloalkyl, aryl, heteroaryl, and  $-\text{CON} R^{15} R^{16}$  or  $-\text{CSN} R^{15} R^{16}$ , in which

$R^{15}$  and  $R^{16}$  independently of one another represent identical or different substituents of the group hydrogen, in each case optionally substituted alkyl or cycloalkyl.